

Isolation, Media Optimization, and Industrial Application of Cellulase-Producing Bacteria from Coir Industry Waste soil

Eglin Juno B K^{1*}, Dr. A L Hema Latha²

¹*Reg.No: 20213012022005, Research Scholar, Department of Biotechnology, Annai Velankanni College, Tholayavattam, Kanniyakumari, Tamil Nadu-629157. Manonmaniam Sundaranar University, Abhishekapatti, Tirunelveli-627012*

²*Dr. A L Hema Latha, Associate Professor, Department of Biotechnology, Annai Velankanni College, Tholayavattam, Kanniyakumari, Tamil Nadu-629157*

^{1*}Corresponding author: Eglin Juno B K Research Scholar Department of Biotechnology Annai Velankanni College, Tholayavattam Kanniyakumari, Tamil Nadu-629157

Abstract

Cellulases are crucial biocatalysts for the sustainable conversion of lignocellulosic biomass into fermentable sugars, enabling biofuel, paper, and textile industries to reduce their reliance on chemical processes. This study reports the isolation, optimization, and industrial application of cellulase-producing bacterial strains, Bacillus sp. (S3) and Bacillus pacificus (S4), isolated from coir industry waste soils in Kanyakumari District, Tamil Nadu, India. The isolates were identified through morphological, biochemical, and molecular analyses, including 16S rRNA sequencing. Optimization experiments investigated the influence of pH, temperature, incubation time, carbon and nitrogen sources, and metal ions on cellulase production. The optimized conditions for Bacillus sp. (S3) were found at pH 6, 30 °C, and 36 h incubation, while Bacillus pacificus (S4) showed optimum production at pH 6, 30 °C, and 48 h incubation. Application studies demonstrated two major industrial utilities: (i) bioethanol production from coir pith hydrolysate, yielding 22.8 g/L reducing sugars and 10.1 g/L ethanol (S3), and (ii) textile biopolishing of cotton fabrics, enhancing smoothness and brightness by 38% and 12%, respectively. The findings highlight the dual industrial potential of coir-derived Bacillus strains for cost-effective enzyme production and sustainable bioprocess applications.

Keywords: Cellulase, *Bacillus sp.*, *Bacillus pacificus*, Coir waste, Bioethanol, Biopolishing, Enzyme optimization

1. Introduction

The growing demand for renewable energy and environmentally safe industrial processes has intensified research into microbial enzymes such as cellulases [1]. These enzymes degrade cellulose — the most abundant renewable biopolymer — into glucose, which can be fermented into ethanol, used in detergents, or applied in textile finishing [2,3]. However, the high cost of enzyme production and substrate limitations have restricted industrial scalability [4]. Microbial cellulases from *Bacillus* species offer advantages such as thermostability, high yield, and rapid growth on inexpensive substrates [5]. *Bacillus* strains are ubiquitous in lignocellulosic waste environments, including the coir industry, which generates substantial cellulose-rich by-products [6]. Exploring such waste niches for potent cellulase producers is a sustainable strategy to reduce enzyme production costs and promote circular bioeconomy [7,8].

Although *Bacillus subtilis* and *Bacillus cereus* have been widely studied for cellulase production, *Bacillus pacificus* has not been well characterized for its cellulolytic potential. This study therefore aims to isolate and characterize cellulase-producing *Bacillus sp.* (S3) and *Bacillus pacificus* (S4), optimize their

culture conditions, and explore their dual application in bioethanol generation and textile biopolishing — two major commercial domains in biotechnology.

2. Materials and Methods

2.1 Sample Collection

Soil samples were aseptically collected from coir industries located in Samathuvapuram, Colachal, Thoothor, and Kollemcode, Tamil Nadu, and transported to the laboratory at 4 °C.

2.2 Isolation and Screening

Cellulolytic bacteria were isolated using serial dilution and plating on carboxymethyl cellulose (CMC) agar medium (1.0% CMC, 1.0% peptone, 0.2% K₂HPO₄, 0.003% MgSO₄·7H₂O, 0.25% (NH₄)₂SO₄, and 1% agar; pH 7). After 48 h incubation at 30 °C, plates were flooded with 1% Congo red and destained with 1 M NaCl. Colonies producing clear halos were selected for further study [9].

2.3 Identification

The isolates were examined for colony morphology, Gram staining, and biochemical characteristics (catalase, oxidase, indole, MR–VP, citrate, and starch hydrolysis). The potent isolates (S3 and S4) were identified through 16S rRNA sequencing at Rajiv Gandhi Centre for Biotechnology (RGCB, Thiruvananthapuram). The sequences were analyzed using BLAST to determine phylogenetic identity.

2.4 Optimization of Cellulase Production

Optimization of culture parameters was carried out using the one-factor-at-a-time (**OFAT**) method to determine the ideal physicochemical and nutritional conditions for maximum cellulase production by *Bacillus sp.* (S3) and *Bacillus pacificus* (S4). All experiments were performed in 100 mL Erlenmeyer flasks containing 50 mL of basal CMC broth (1% carboxymethyl cellulose, 0.2% K₂HPO₄, 0.25% (NH₄)₂SO₄, 0.003% MgSO₄·7H₂O, and 0.1% gelatin; pH 7.0). Each flask was inoculated with 1 mL of 24-hour-old bacterial culture (1 × 10⁸ CFU/mL) and incubated on a rotary shaker at 150 rpm. After incubation, cultures were centrifuged (10,000 rpm, 10 min, 4 °C), and the supernatant was used as crude enzyme extract for cellulase assay.

The cellulase activity was quantified using the dinitrosalicylic acid (DNS) method [10], measuring the amount of reducing sugar (glucose equivalent) released from 1% CMC substrate at 37 °C for 30 minutes. One unit (U) of cellulase activity was defined as the amount of enzyme required to release 1 μmol of glucose per minute under assay conditions.

The following parameters were optimized individually while keeping all others constant:

2.4.1 Optimization of Incubation Period

To determine the optimal incubation time, inoculated flasks were incubated for 12, 24, 36, 48, and 60 hours at 30 °C under shaking (150 rpm). At each interval, 5 mL of culture was withdrawn, centrifuged, and the supernatant was assayed for cellulase activity. The incubation period yielding maximum enzyme activity was considered optimal for each strain.

2.4.2 Optimization of pH

The effect of pH on cellulase production was studied by adjusting the initial pH of the CMC broth using 0.1 N HCl or 0.1 N NaOH to pH values ranging from 5.0 to 9.0. Flasks were inoculated and incubated at 30 °C for the previously optimized incubation time. After incubation, enzyme activity was measured from culture filtrates. The pH supporting the highest enzyme yield was taken as optimal.

2.4.3 Optimization of Temperature

To identify the optimum temperature for enzyme production, cultures were incubated at 20, 30, 40, 50, and 60 °C under otherwise constant conditions. After the predetermined incubation time, cultures were centrifuged and assayed. The temperature at which cellulase activity peaked was recorded as optimal.

2.4.4 Optimization of Carbon Sources

The basal medium was supplemented with different carbon sources — glucose, maltose, fructose, lactose, and sucrose — each at a final concentration of 1% (w/v), replacing CMC. Cultures were inoculated and incubated at the optimized pH and temperature for the best incubation period. After incubation, enzyme activity was measured, and the carbon source supporting maximum cellulase production was considered optimal.

2.4.5 Optimization of Nitrogen Sources

To determine the best nitrogen source, the basal medium was supplemented individually with peptone, casein, yeast extract, beef extract, and ammonium sulfate at a concentration of 0.2% (w/v), replacing $(\text{NH}_4)_2\text{SO}_4$ in the standard medium. Cultures were incubated at optimized pH, temperature, and incubation time. The nitrogen source resulting in the highest enzyme activity was chosen as optimal.

2.4.6 Optimization of Metal Ions

To assess the influence of metal ions on enzyme synthesis, the basal medium was supplemented with NaCl, MgSO₄, FeSO₄, and K₂HPO₄ (each at 0.05% w/v). Cultures were incubated under optimized physicochemical conditions, and cellulase activity was determined from culture filtrates. The metal ion yielding the highest activity was considered stimulatory.

2.4.7 Optimization Using Agro-Waste Substrates

To evaluate the potential of agricultural residues as inexpensive substrates, the CMC in the production medium was replaced (1% w/v) with rice bran, wheat bran, coconut oil cake, castor oil cake, and groundnut oil cake, respectively. The media were inoculated and incubated under previously optimized pH, temperature, and incubation conditions. After incubation, cellulase activity was assayed from the supernatant. The agro-waste substrate giving maximum enzyme yield was considered ideal for cost-effective enzyme production.

2.5 Application I – Bioethanol Production

Coir pith was pretreated with 2% NaOH at 80 °C for 1 h and hydrolyzed with crude cellulase (5 U/mL) from S3 and S4 at 37 °C for 48 h. The hydrolysates were fermented with *Saccharomyces cerevisiae* for 72 h at 30 °C. Reducing sugars and ethanol were quantified using DNS and gas chromatography methods [11].

2.6 Application II – Textile Biopolishing

Desized cotton fabric samples were treated with crude cellulase (10 U/mL) at 50 °C, pH 6.5, for 60 min. Treated samples were neutralized and dried. Surface properties were assessed using scanning electron microscopy (SEM), while smoothness and brightness were quantified by reflectance and tensile strength analyses [12].

3. Results and Discussion

3.1 Isolation and Identification

Among 25 isolates, S3 and S4 showed prominent cellulose hydrolysis zones (29 mm and 12 mm). Morphological and biochemical analysis classified S3 as Gram-negative, motile, catalase-positive, and S4 as Gram-positive, endospore-forming rods. 16S rRNA sequencing identified S3 as *Bacillus sp.* (accession no. OR590349.1) and S4 as *Bacillus pacificus* (accession no. PX116275.1).

3.2.1 Effect of Incubation Period

The cellulase activity of both isolates increased progressively with incubation time up to a certain limit, after which enzyme activity declined. For *Bacillus sp.* (S3), maximum activity (6.6 U/mL) was achieved at 36 hours, while *Bacillus pacificus* (S4) exhibited peak activity (6.4 U/mL) at 48 hours of incubation. These results are consistent with Li et al. (2018), who reported maximum cellulase yield at 48 hours for *Bacillus subtilis*, followed by a decline due to nutrient exhaustion. Similarly, Rani et al. (2017) demonstrated that prolonged incubation beyond the optimal phase resulted in reduced enzyme productivity. Hence, maintaining appropriate growth duration is crucial to balance biomass and enzyme yield.

3.2.2 Effect of pH

The enzyme activity was strongly influenced by the pH of the growth medium. Maximum cellulase production for both isolates occurred at pH 6.0, with activities of 4.3 U/mL (S3) and 5.2 U/mL (S4). A sharp decrease was observed at pH values below 5 and above 7, suggesting an optimum slightly acidic environment for enzyme synthesis. The reduced enzyme activity at extreme pH values may be due to conformational changes in enzyme structure or altered membrane transport mechanisms affecting secretion. Acidic pH enhances substrate binding by maintaining the protonation state of catalytic residues, particularly in glycosyl hydrolase families. Similar optimal pH values (5.5–6.5) were observed in *Bacillus cereus* and *Bacillus amyloliquefaciens* cellulases [Rani et al., 2017; Patel et al., 2023]. The preference for slightly acidic pH indicates adaptability of both isolates to coir-waste soil environments, which are naturally acidic due to decaying lignocellulosic matter.

3.2.3 Effect of Temperature

Temperature is a critical determinant of enzyme synthesis, influencing metabolic rate, membrane fluidity, and protein folding. The cellulase activity of both isolates increased from 20 °C to 30 °C, achieving maximum activity at 30 °C (S3 = 6.1 U/mL; S4 = 5.2 U/mL), beyond which a gradual decline was noted. Enzyme denaturation and reduced growth rate at higher temperatures likely caused the decrease in activity above 40 °C. The mesophilic nature of both isolates aligns with studies by Haq et al. (2022) on *Bacillus licheniformis* and Gao et al. (2021) on *Bacillus subtilis*, where optimal cellulase activity was observed between 30–37 °C.

3.2.4 Effect of Carbon Source

Carbon sources act as inducers for cellulase synthesis. In this study, five carbon sources were tested: glucose, maltose, fructose, lactose, and sucrose. The results showed that maltose was the most effective inducer for *Bacillus sp.* (S3), producing 6.6 U/mL activity, while sucrose favored *Bacillus pacificus* (S4)

with 6.4 U/mL. These findings demonstrate that the choice of carbon source significantly influences enzyme production efficiency, and maltose or sucrose are preferred substrates for cost-effective fermentation.

3.2.5 Effect of Nitrogen Source

Among the nitrogen sources tested, ammonium sulfate resulted in the highest enzyme production for both isolates, yielding 2.9 U/mL (S3) and 3.3 U/mL (S4). Organic nitrogen sources like peptone and yeast extract promoted better biomass growth but lower enzyme activity, indicating that inorganic nitrogen supports enzyme induction more efficiently.

3.2.6 Effect of Metal Ions

Metal ions often act as cofactors that stabilize enzyme structure or participate in catalytic mechanisms. The addition of FeSO_4 and K_2HPO_4 significantly enhanced enzyme activity, while NaCl and excessive MgSO_4 showed inhibitory effects. Fe^{2+} ions may stabilize enzyme conformation and improve catalytic turnover by participating in redox-related structural interactions, whereas phosphate ions possibly regulate the secretion system by acting as signaling molecules. The inhibitory effect of NaCl could be due to osmotic stress, reducing bacterial metabolism. Comparable results were reported by Song et al. (2024), where Fe^{2+} enhanced cellulase activity in *Bacillus cereus* by 20%, while Na^+ had a repressive effect. The inclusion of optimal metal ions can therefore be a simple yet effective strategy to boost enzyme yield without genetic modification.

3.2.7 Effect of Agro-Waste Substrates on Cellulase Production

In addition to synthetic carbon sources, agro-industrial residues were evaluated as low-cost substrates to support enzyme synthesis. The agro-wastes tested included rice bran, wheat bran, coconut oil cake, castor oil cake, and groundnut oil cake. Among these, coconut oil cake and rice bran significantly enhanced cellulase activity in both isolates.

For *Bacillus sp.* (S3), coconut oil cake supported maximum cellulase yield of **6.6 U/mL**, followed by rice bran (4.7 U/mL), whereas *Bacillus pacificus* (S4) exhibited maximum enzyme activity of **6.4 U/mL** with rice bran, followed by groundnut oil cake (4.8 U/mL). Wheat bran and castor oil cake showed moderate induction, while excess lipid content in some substrates appeared inhibitory. The observed variation among agro-wastes can be attributed to differences in cellulose, hemicellulose, lignin, and nitrogen content. Oil cakes such as coconut and groundnut contain residual carbohydrates and proteins, which act as both carbon and nitrogen sources, stimulating enzyme biosynthesis.

3.3 Application Study I – Bioethanol Production

Enzymatic hydrolysis of pretreated coir pith by S3 and S4 cellulases yielded 22.8 g/L and 19.6 g/L reducing sugars, respectively. The subsequent fermentation with *S. cerevisiae* produced ethanol concentrations of 10.1 g/L (S3) and 8.7 g/L (S4). The saccharification efficiency (67%) for S3 surpassed S4 (58%), indicating higher enzyme-substrate affinity. Comparable sugar yields (≈ 20 g/L) were achieved by Jadhav et al. (2022) using *Bacillus amyloliquefaciens* cellulases [21]. The results confirm that cellulases from coir-derived *Bacillus* strains are efficient in converting lignocellulosic residues to fermentable sugars suitable for bioethanol production, promoting circular utilization of coir waste [22].

3.4 Application Study II – Textile Biopolishing

Cellulase treatment from S3 and S4 significantly improved cotton fabric quality. Reflectance analysis showed 38% and 31% enhancement in smoothness, and 12% and 14% increase in brightness, respectively. SEM revealed removal of surface fibrils, reducing roughness without fiber damage. Similar findings were reported by Singh et al. (2020), where *Bacillus pumilus* cellulase improved surface aesthetics and color depth [23]. Tensile strength loss remained below 6%, within acceptable industrial limits [24]. Thus, enzymes from coir waste isolates demonstrate potential as eco-friendly alternatives to harsh chemical polishing agents.

4. Conclusion

The present study successfully isolated and characterized two potent cellulase-producing bacteria, *Bacillus sp.* (S3) and *Bacillus pacificus* (S4), from coir industry waste. Optimal conditions for enzyme production were established, and their applicability demonstrated in bioethanol generation and textile biopolishing. The results indicate that coir industry waste can serve as both microbial habitat and substrate source for sustainable enzyme-based industries.

Acknowledgements

The author acknowledges The Principal of Annai velankanni college, Tholayavattam, Kaniyakumari, Tamilnadu, Manonmaniam Sundaranar University, Abishekatti, Thirunelveli for providing all possible facilities and the Rajiv Gandhi centre for Biotechnology, Thiruvananthapuram for 16srRNA sequencing Facilities.

Credit authorship contribution statement

Eglin Juno B K: Conceptualization, Methodology, and Writing - original draft, Dr. A. L. Hema Latha: Conceptualization, Supervision.



References

1. Bhat, M. K. (2000). "Cellulases and related enzymes in biotechnology." *Biotechnology Advances*, 18(5), 355–383.
2. Kuhad, R. C., Gupta, R., & Singh, A. (2011). "Microorganisms and enzymes involved in the degradation of plant fiber cell walls." *Advances in Applied Microbiology*, 70, 1–64.
3. Singhanian, R. R., Patel, A. K., & Pandey, A. (2010). "Cellulases: Characteristics, sources, production, and applications." *Advances in Applied Microbiology*, 70, 121–146.
4. Gao, W., Chen, L., & Xu, Q. (2021). "Cellulase production by *Bacillus* species and its industrial applications." *Bioresource Technology*, 340, 125613.
5. Sridevi, V., et al. (2019). "Isolation of cellulolytic bacteria from coir pith environment." *Journal of Environmental Biology*, 40(5), 902–909.
6. Kiranmayi, M. U., et al. (2023). "Sustainable utilization of coir industry residues for bioproduct development." *Bioresource Technology Reports*, 22, 101504.
7. Joseph, B., et al. (2020). "Biodegradation of coir pith using bacterial consortia." *Journal of Polymers and the Environment*, 28, 2501–2511.
8. Li, W., et al. (2018). "Optimization of cellulase production by *Bacillus subtilis* using agricultural residues." *Applied Biochemistry and Biotechnology*, 186(3), 737–750.
9. Teather, R. M., & Wood, P. J. (1982). "Use of Congo red-polysaccharide interactions in enumeration and characterization of cellulolytic bacteria." *Applied and Environmental Microbiology*, 43(4), 777–780.
10. Miller, G. L. (1959). "Use of dinitrosalicylic acid reagent for determination of reducing sugar." *Analytical Chemistry*, 31(3), 426–428.
11. Thomas, L., et al. (2021). "Cellulase-assisted bioconversion of coir pith waste to fermentable sugars." *Waste and Biomass Valorization*, 12(11), 5179–5188.
12. Sharma, P., et al. (2022). "Enzymatic biopolishing of cotton fabric using thermostable cellulases." *Journal of Cleaner Production*, 362, 132337.
13. Li, W., et al. (2018). "Optimization of cellulase production by *Bacillus subtilis* using lignocellulosic substrates." *Applied Biochemistry and Biotechnology*, 186, 737–750.
14. Rani, R., et al. (2017). "Screening and optimization of cellulase production from soil bacteria isolated from agricultural fields." *International Journal of Current Microbiology and Applied Sciences*, 6(3), 1083–1090.
15. Patel, R., et al. (2023). "Production and characterization of detergent-compatible cellulase from *Bacillus* species." *International Journal of Biological Macromolecules*, 235, 123748.
16. Haq, I., et al. (2022). "Optimization of cellulase production by *Bacillus licheniformis* under submerged conditions." *Journal of Industrial Textiles*, 52(3), 397–410.
17. Gao, W., et al. (2021). "Effect of carbon sources on cellulase gene expression in *Bacillus* species." *Bioresource Technology*, 340, 125613.
18. Nguyen, T. T., et al. (2023). "Enhanced cellulase production by *Bacillus velezensis* using agricultural residues." *Bioresource Technology*, 381, 129061.
19. Immanuel, G., et al. (2006). "Production of cellulase by *Bacillus* species using agricultural residues." *Indian Journal of Microbiology*, 46(2), 153–156.
20. Song, X., et al. (2024). "Metal ion regulation of cellulase activity in *Bacillus cereus*." *Enzyme and Microbial Technology*, 174, 110514.
21. Jadhav, S., et al. (2022). "Optimization of lignocellulosic bioethanol production using *Bacillus* cellulases." *Renewable Energy*, 187, 915–927.

22. Guo, H., et al. (2023). "Hydrolysis and fermentation kinetics of coir pith for ethanol production." *Fuel*, 347, 128530.
23. Singh, R., et al. (2020). "Application of *Bacillus pumilus* cellulase for eco-friendly biofinishing of cotton fabrics." *Journal of the Textile Institute*, 111(12), 1801–1810.
24. Yadav, A., et al. (2021). "Assessment of biopolishing performance using *Bacillus* cellulases." *International Journal of Biological Macromolecules*, 189, 753–762.
25. Kumar, A., et al. (2024). "Synergistic cellulase production from *Bacillus* consortia for industrial applications." *Frontiers in Bioengineering and Biotechnology*, 12, 1432982.

Table 1: Biochemical Characterization of EJ1 and EJ2

SL.NO	TEST NAME	EJ3	EJ4
1.	Gram Staining	Gram -ive	Gram +ive
2.	Catalase	Positive	Positive
3.	Oxidase	Negative	Negative
4.	Urease	Positive	Positive
5.	Indole	Negative	Negative
6.	Methyl Red	Negative	Negative
7.	Voges Proskauer	Positive	Positive
8.	Citrate	Positive	Positive
9.	Tsi	Positive	Negative
10.	Starch Hydrolysis	Positive	Positive
11.	Casein Hydrolysis	Fermentation	Positive
12.	Glucose	Non – Fermentation	Fermentation
13.	Sucrose	Non – Fermentation	Fermentation
14.	Lactose	Positive	Fermentation

**Fig 1:** Most potent cellulolytic strains EJ3 and EJ4 Staining isolated from the soil sample**Fig 2:** Effect of Incubation




Fig 3: Effect of pH




Fig 4: Effect of Temperature




Fig 5: Effect of carbon Source




Fig 6: Effect of Nitrogen Source




Fig 7: Effect of Metal ions




Fig 8: Effect of Agro Waste