

FUZZY TOPOLOGICAL SPACE WITH AN BOOLEAN ALGEBRA

Dr. K. Balasubramaniyan¹ *C.Dilly Rani²

¹Assistant Professor, Department of Mathematics, Annamalai University, Chidambaram.
Deputed to Arignar Anna Government Arts College,
Vadachennimalai-636121, Attur, Tamilnadu, India.

² J.J.College of Engineering and Technology, Tiruchirappalli-600009, Tamilnadu, India.

Abstract: In this paper, we introduce the fuzzy topology by involving the Boolean algebraic structure on it and the notion of Boolean algebraic fuzzy topological spaces. Fuzzy set was introduced by L.A. Zadeh as a generalization of the notion of classical set or crisp set. Fuzzy topological spaces were introduced by C.L. Chang and studied by many eminent authors like C.K.Wong. A.Rosenfeld applied the notion of fuzzy set to algebra and introduced fuzzy subgroup of a group. Shaoquan Sun introduced the notion of fuzzy Boolean subalgebra in a Boolean algebra.

Keywords: Boolean algebraic fuzzy topological space (BAFTS), Fuzzy topological space(FTS), Fuzzy Boolean subalgebra(FBSA),

1. Introduction

L.A Zadeh's [8] introduced the fundamental concept of fuzzy sets. The study of fuzzy topological spaces and fuzzy Boolean algebra are the two main branches of such mathematics. General topology and Boolean algebra can be regarded as a special case of fuzzy topology and fuzzy Boolean algebra respectively, such as membership function takes the values 0 and 1 only. Some basic properties of fuzzy topology are discussed in [1,10] and related to fuzzy subalgebra of Boolean algebra are discussed in [5, 6]. It is remarked here that Boolean algebraic fuzzy topological spaces are different from fuzzy topological Boolean algebras, as introduced by A. Parvathi and K.N. Meenakshi in [3]

2. Preliminaries

In this section, we list some basic concepts and well known results on fuzzy topology and fuzzy Boolean algebra for the sake of completeness of the topic under study. Throughout the paper, $\emptyset_{\sim}$ and $X_{\sim}$ denotes the constant fuzzy sets on X defined by $\emptyset_{\sim}(x) = 0; \forall x \in X$ and $X_{\sim}(x) = 1; \forall x \in X$.

Definition(2.1): Let X be an non empty set and I^X denotes the set of all fuzzy subsets of X , i.e., the set of all mappings $\mu: X \rightarrow I$. Then a subfamily $\delta \subseteq I^X$ is said to be a fuzzy topology on X , if it satisfies the following:

- (i) $\emptyset \sim, X \sim \in \delta$;
- (ii) If $\{\mu_i : i \in \Lambda\} \subseteq \delta$, then $\bigcup_{i \in \Lambda} \mu_i \in \delta$;
- (iii) If $\mu, \nu \in \delta$ be any two members, then $\mu \cap \nu \in \delta$.

If δ is a fuzzy topology on X , then the pair (X, δ) is called fuzzy topological space (FTS). The members of δ are called δ -open sets. A fuzzy set μ of X is said to be δ -closed in (X, δ) if and only if μ^c is δ -open set in (X, δ) . When $\delta = I^X$, then (X, δ) is called a discrete fuzzy topological space and when $\delta = \{\emptyset \sim, X \sim\}$, then (X, δ) is called an indiscrete fuzzy topological space.

Definition (2.2): Let X and Y be two non-empty sets and $f: X \rightarrow Y$ be a mapping. Let μ and ν be fuzzy sets of X and Y respectively. Then the image of μ under the map f is denoted by $f(\mu)$ and is defined as defined as $xf(\mu)(y) = \{Sup \{\mu(x) : x \in f^{-1}(y)\} : I | f^{-1}(y) \neq \emptyset\}$. Also the pre image of ν under f is denoted by $f^{-1}(\nu)$ and is defined as $f^{-1}(\nu)(x) = \nu(f(x))$; $\forall x \in X$.

Remark (2.3) : Note that $\mu(x) \leq f(\mu)(f(x))$; $\forall x \in X$, and equality hold when f is bijective.

Definition (2.4)[1]: Let (X_1, δ_1) and (Y_2, δ_2) be FTSs. Then the function $f: (X_1, \delta_1) \rightarrow (Y_2, \delta_2)$ is

- (i) fuzzy continuous: if and only if for every $\nu \in \delta_2$; $f^{-1}(\nu) \in \delta_1$
- (ii) fuzzy open: if and only if for every $\mu \in \delta_1 \Rightarrow f(\mu) \in \delta_2$.
- (iii) fuzzy closed: if and only if image of every δ_1 -closed set is δ_2 -closed set.
- (iv) fuzzy homomorphism: if and only if f is bijective, fuzzy continuous and fuzzy open (or fuzzy closed).

Definition (2.5)[10]: Let ν be a fuzzy set in X and δ is a fuzzy topology on X . Then the induced fuzzy topology on ν is the family of subsets of ν which are the intersections with ν of δ -open fuzzy sets in X . The induced fuzzy topology is denoted by δ_ν i.e., $\delta_\nu = \{\mu \cap \nu : \forall \mu \in \delta\}$, and the pair (ν, δ_ν) is called a fuzzy subspace of (X, δ) .

Definition (2.6)[10]: For any $\lambda \in (0, 1]$ and $x \in X$, a fuzzy set x_λ in X is called a fuzzy point in X if

$$x_\lambda(y) = \begin{cases} \lambda & \text{if } y = x, \forall y \in X \\ 0 & \text{otherwise} \end{cases}$$

The fuzzy point x_λ is said to belongs to the fuzzy set μ , denoted by $x_\lambda \in \mu$, iff $\mu(x) \geq \lambda$. In particular, $x_\lambda \leq y_\sigma \Leftrightarrow x = y$ and $\lambda \leq \sigma$. The characteristic mapping of a subset A of X is denoted by χ_A and is defined as

$$\chi_A(x) = \begin{cases} 1, & \text{if } x \in A \\ 0, & \text{otherwise} \end{cases}$$

Obviously, a characteristic function χ_A is also a fuzzy set on X and for any non-empty subsets A and B of X , we have $A \subseteq B$ if and only if $\chi_A \subseteq \chi_B$.

Definition(2.7)[5]: Let $(B, +, *, ', 0_B, 1_B)$ be a Boolean algebra and $\mu \in I^B$ be a fuzzy set, then μ is called a fuzzy Boolean subalgebra (FBSA) of B if for all $x, y \in B$

- (i) $\mu(x+y) \geq \min\{\mu(x), \mu(y)\};$
- (ii) $\mu(x*y) \geq \min\{\mu(x), \mu(y)\};$
- (iii) $\mu(x') = \mu(x).$

Lemma (2.8): For any fuzzy Boolean subalgebra μ of Boolean algebra $(B, +, *, ', 0_B, 1_B)$ the following holds

- (i) $\mu(1_B) \geq \mu(x), \forall x \in B$
- (ii) $\mu(0_B) \geq \mu(x), \forall x \in B$
- (iii) $\mu(0_B) = \mu(1_B)$

Proof:(i) Since $\mu(1_B) = \mu(x+x') \geq \min\{\mu(x), \mu(x')\} = \min\{\mu(x), \mu(x)\} = \mu(x), \forall x \in B$

(ii) Also, $\mu(0_B) = \mu(xx') \geq \min\{\mu(x), \mu(x')\} = \min\{\mu(x), \mu(x)\} = \mu(x), \forall x \in B$

(iii) From (i) and (ii), we get $\mu(1_B) \geq \mu(0_B)$ and $\mu(0_B) \geq \mu(1_B)$ and so $\mu(0_B) = \mu(1_B)$.

The following results are

- Result(2.9):**(i) If H is a subset of a Boolean algebra B , then χ_H is a fuzzy Boolean subalgebra of B if and only if H is a Boolean subalgebra of B .
- (ii) All constant fuzzy sets of a Boolean algebra B are fuzzy Boolean subalgebra of B .

3. Fuzzy Topology with Boolean Algebraic

In this section, we will introduce Boolean algebraic fuzzy topology and give basic properties of this structure. We also discuss similarities with and difference from other fuzzy topological spaces.

Definition (3.1): Let $(B, +, *, ', 0_B, 1_B)$ be a Boolean algebra and I^B denotes the set of all fuzzy subsets of B . Then the collection $T = \{\mu: \mu \in I^B \text{ such that } \mu(x) = \mu(x') \text{ for all } x \in B\}$ is a fuzzy topology on B and B with this topology is called **Boolean algebraic fuzzy topological space (BAFTS)**.

It is denoted by (B, T) .

Proof. Since $\emptyset_{\sim}$ and $B_{\sim}$ are constant fuzzy sets on B such that $\emptyset_{\sim}(x) = \emptyset_{\sim}(x') = 0$, and $B_{\sim}(x) = B_{\sim}(x') = 1$, for all $x \in B$. Therefore, $\emptyset_{\sim} \in T$ and $B_{\sim} \in T$.

Next, let $\mu, \nu \in T$ be any two members and $x \in B$ be any element, then

$(\mu \cap \nu)(x) = \min\{\mu(x), \nu(x)\} = \min\{\mu(x'), \nu(x')\} = (\mu \cap \nu)(x')$. Thus $\mu \cap \nu \in T$.

Further, let $\{\mu_i : i \in \Lambda\} \subseteq T$.

$$\left(\bigcup_{i \in \Lambda} \mu_i\right)(x) = \sup_{i \in \Lambda} \{\mu_i(x_i)\} = \sup \{\mu_i(x')\} = \left(\bigcup_{i \in \Lambda} \mu_i\right)(x').$$

So that $(\bigcup_{i \in \Lambda} \mu_i) \in T$.

Hence T is a fuzzy topology on B .

Remark (3.2): A fuzzy set μ of a BAFTS (B, T) is called T -closed if $\mu^c \in T$. It is easily to verify that $\emptyset$ and B are T -closed and if μ, ν are T -closed, then $\mu \cup \nu$ is also T -closed and if $\{\mu_i : i \in \Lambda\}$ be an arbitrary family of T -closed sets, then $\bigcap_{i \in \Lambda} \mu_i$ is also T -closed set.

Theorem (3.3) : A fuzzy set μ of the BAFTS B is T -closed if and only if μ is T -open.

Proof: Let μ be T -open set, then $\mu(x) = \mu(x')$ for all $x \in B$.

Now, $\mu^c(x) = 1 - \mu(x) = 1 - \mu(x') = \mu^c(x')$.

So μ^c is T -open i.e., μ is T -closed.

Converse is also true.

Example(3.4): Let $B = \{0, a, b, 1\}$ be the diamond Boolean algebra, where the three operations $+$, $*$ and $'$ are defined as $x + y = \sup\{x, y\}$ and $x * y = \inf\{x, y\}$ such that $0' = 1, 1' = 0, a' = b$ and $b' = a$.

Let $\delta = I^B$ be the discrete fuzzy topology on B . We show that (B, δ) is not a BAFTS.

Let $H = \{0, a, 1\}$ be a subset of B . Define a fuzzy set μ on B as follows:

$$\mu(x) = \begin{cases} \lambda & \text{if } x \in H \\ 0 & \text{if } x \in B \setminus H \end{cases}, \text{ where } \lambda \in (0, 1]$$

Then $\mu(a) = \lambda$, but $\mu(a') = \mu(b) = 0$ i.e., $\mu(a) \neq \mu(a')$, so $\mu \notin T$, where T is the BAFT on B . Hence δ is not a BAFT on B .

Now, we show that under certain condition on B , a BAFT S is also a discrete FTS.

Theorem(3.5): Let (B, T) be a BAFTS such that $x' = x, \forall x \in B$. Then (B, T) is a discrete FTS.

Proof: Let $\mu \in I^B$ and $x \in B$ be any element. Then $\mu(x) = \mu(x') \Rightarrow \mu \in T$. So $I^B \subseteq T$. Hence T is a discrete fuzzy topology on B .

Note (3.6): We notice that the Theorem (3.5) holds only in the case when the Boolean algebra B is trivial one. This implies that there does not exist non-trivial Boolean algebra B for which the BAFTS is a discrete fuzzy topological space.

Remark (3.7): (i) If B is a Boolean algebra such that there exist atleast one element x , such that $x \notin \{0B, 1B\}$,

then the BAFTS on B is not a discrete FTS.

(ii) If $|B| \geq 2$, then the BAFTS (B, T) is not an indiscrete FTS, i.e., there exist $\mu \in I^B$, which is different from $\emptyset$ and B such that $\mu \in T$.

Theorem (3.8): Every fuzzy Boolean subalgebra of a Boolean algebraic fuzzy topological space B is T -open set.

Proof: Let κ be fuzzy Boolean subalgebra of a Boolean algebra B . Then by Definition(2.8), we have $\kappa(x') = \kappa(x)$, $\forall x \in B$ i.e., $\kappa \in T$. Hence κ is T -open set.

Remark (3.9): Converse of Theorem (3.7) is not true i.e., T -open set does not need to be fuzzy Boolean subalgebra of B .

Example (3.10): Consider $B = \{0B, a, b, 1B\}$ be the diamond Boolean algebra as defined in Example(3.4).

Let $H = \{a, b\}$ be a subset of B . Define a fuzzy set μ on B as follows:

$$\mu(x) = \begin{cases} \lambda & \text{if } x \in H \\ 0 & \text{if } x \in B \setminus H \end{cases}, \text{ where } \lambda \in (0, 1]$$

Let T be the BAFT defined on B . If $\mu \in T$ however, μ is an fuzzy Boolean subalgebra of B , for

$\mu(a+b) = \mu(1B) = 0 \geq \lambda \neq \min\{\mu(a), \mu(b)\}$. If T is a open set in BAFTS (B, T) is a fuzzy Boolean subalgebra of B .

Lemma(3.11): Let μ be a fuzzy Boolean subalgebra of the Boolean algebra B , then $(xy)_\gamma \in \mu$ and $(x+y)_\gamma \in \mu$ where $\gamma = \min\{\alpha, \beta\} \forall x_\alpha, y_\beta \in \mu$

Proof. Let μ be a fuzzy Boolean subalgebra of B and $x_\alpha, y_\beta \in \mu$ then, we have

$$\mu(x) \geq \alpha \text{ and } \mu(y) \geq \beta \therefore \min\{\mu(x), \mu(y)\} \geq \min\{\alpha, \beta\} = \gamma.$$

As μ is fuzzy Boolean subalgebra of B

$$\therefore \mu(xy) \geq \min\{\mu(x), \mu(y)\} \geq \min\{\alpha, \beta\} = \gamma \text{ and } \mu(x+y) \geq \min\{\mu(x), \mu(y)\} \geq \min\{\alpha, \beta\} = \gamma.$$

This implies that $(xy)_\gamma \in \mu$ and $(x+y)_\gamma \in \mu$.

Theorem (3.12): Let μ be T-open set in BAFTS (B,T), then μ is a fuzzy Boolean subalgebra of B if and only if $\forall x, y \in B$, we have $(xy)_{\gamma} \in \mu$ and $(x+y)_{\gamma} \in \mu$, where $\gamma = \min\{\alpha, \beta\}$.

Proof: When $\mu \in T$ and μ is a fuzzy Boolean subalgebra of B, then the result follows by Lemma (3.11)

Conversely, let μ be T-open set such that $\forall x, y \in B$, we have $(xy)_{\gamma} \in \mu$ and $(x+y)_{\gamma} \in \mu$, where $\gamma = \min\{\alpha, \beta\}$

To show that μ is a fuzzy Boolean subalgebra of B.

Let $x, y \in B$ be any element and suppose that $\mu(x) = \alpha$ and $\mu(y) = \beta$. Suppose

that $\alpha \leq \beta$, then $\min\{\alpha, \beta\} = \alpha$.

Now, $\mu(x) = \alpha$ and $\mu(y) = \beta \geq \alpha$. Then $x_{\alpha}, y_{\alpha} \in \mu \Rightarrow (xy)_{\alpha} \in \mu$ and $(x+y)_{\alpha} \in \mu$

i.e., $\mu(xy) \geq \alpha = \min\{\alpha, \beta\} = \min\{\mu(x), \mu(y)\}$ and $\mu(x+y) \geq \alpha = \min\{\mu(x), \mu(y)\}$.

If $\beta \leq \alpha$ is similar. Thus in both the cases, we find that

$\mu(xy) \geq \min\{\mu(x), \mu(y)\}$ and $\mu(x+y) \geq \min\{\mu(x), \mu(y)\}$.

As $\mu \in T$, therefore, $\mu(x) = \mu(x')$ for all $x \in B$.

Hence μ is fuzzy Boolean subalgebra of B.

Theorem (3.13): Let H be a Boolean subalgebra of a BAFTS(B,T), then the relative fuzzy topology T_H on H is a BAFT on H.

Proof: The subset H of B (in the ordinary sense) has a characteristic function χ_H such that

$$\chi_H(x) = \begin{cases} 1, & \text{if } x \in H \\ 0, & \text{if } x \notin H \end{cases}$$

Clearly, $\chi_H(x) = 1, \forall x \in H$ is a constant fuzzy set on H.

Let $T_H = \{\mu \cap \chi_H : \mu \in T\}$. Then we claim that T_H is a BAFT on H.

For, choose $\emptyset \sim \in T$, then as $(\emptyset \sim \cap \chi_H)(x) = \min\{\emptyset \sim(x), \chi_H(x)\} = \emptyset \sim(x)$, so we have $\emptyset \sim \in T_H$.

To show that $H \sim \in T_H$, choose χ_B the characteristic function for B, then

$$(\chi_B \cap \chi_H)(x) = \min\{\chi_B(x), \chi_H(x)\} = \chi_H(x), \text{ so we have } H \sim \in T_H.$$

Further, let $\{\mu_i \cap \chi_H : i \in \Lambda\} \subseteq T_H$. Then, $\bigcup_{i \in \Lambda} (\mu_i \cap \chi_H) = \bigcup_{i \in \Lambda} \mu_i \cap \chi_H$.

Also, let $\mu_1 \cap \chi_H$ and $\mu_2 \cap \chi_H$ be any two members of T_H .

Then, $(\mu_1 \cap \chi_H) \cap (\mu_2 \cap \chi_H) = (\mu_1 \cap \mu_2) \cap \chi_H \in T_H$. Hence T_H is a fuzzy topology on H.

Further, let T_1 is a BAFT on H, then we show that $T_1 = T_H$.

Let $\mu \cap \chi_H \in T_H$, then $(\mu \cap \chi_H)(x) = \min\{\mu(x), \chi_H(x)\} = \min\{\mu(x'), \chi_H(x')\} = (\mu \cap \chi_H)(x')$.

So, $\mu \cap \chi_H \in T_1$, i.e., $T_H \subseteq T_1$.

Further, let $v \in T_1$ and so $v(x) = v(x'), \forall x \in B$.

Also, $v = v \cap \chi_H \in T_H$. Therefore, $T_1 \subseteq T_H$ and so $T_1 = T_H$.

Hence relative fuzzy topology on H is the BAFT on H .

4. Functions on Fuzzy Topological Spaces with Boolean Algebraic

In this section, we study the behavior of Boolean algebraic fuzzy topology under different Boolean homomorphism (isomorphism) and also under fuzzy continuity from one Boolean algebraic fuzzy topological space to another Boolean algebraic fuzzy topological space.

Theorem (4.1): Let $f: B_1 \rightarrow B_2$ be a Boolean isomorphism and T_1, T_2 be BAFTs on Boolean algebra B_1 and B_2 respectively. Then f is a fuzzy homomorphism between BAFTS (B_1, T_1) and (B_2, T_2) .

Proof: Let $f: B_1 \rightarrow B_2$ be a Boolean isomorphism, where B_1 and B_2 are Boolean algebras. We show that f is a fuzzy homeomorphism. For this, we need to show that f is a fuzzy continuous and fuzzy open map. For fuzzy continuity: let v be any T_2 - open set and $x \in B_1$ be any element. Then

$$(f^{-1}(v))(x) = v(f(x)) = v[\{f(x)\}'] = v(f(x')) = (f^{-1}(v))(x').$$

So $f^{-1}(v)$ is T_1 -open set. Hence f is fuzzy continuous.

For fuzzy open mapping: let μ be any T_1 -open set and let $y \in B_2$ be any element, then unique $x \in B_1$ such that $f(x) = y$.

$$\text{Now, } f(\mu)(y) = f(\mu)(f(x)) = \mu(x) = \mu(x') = f(\mu)(f(x')) = f(\mu)(f(x')) = f(\mu)(y').$$

i.e., $f(\mu) \in T_2$. Therefore, f is fuzzy open mapping. Hence f is a fuzzy homeomorphism.

Theorem(4.2): Let $f: (B, T) \rightarrow (B, T)$ be a mapping from a BAFTS (B, T) on to itself defined by $f(x) = x'$; $\forall x \in B$. Then f is bijective and fuzzy continuous.

Proof: Clearly, f is one-one and onto. Let $v \in T$ be any open set. Then we show that $f^{-1}(v) \in T$.

Let $x \in B$ be any element $\Rightarrow f(x) = x'$.

$$\text{Now, } f^{-1}(v)(x) = v(f(x)) = v(x') = v(x) = v(f(x')) = f^{-1}(v)(x')$$

So, $f^{-1}(v) \in T$. Hence f is fuzzy continuous.

Theorem(4.3): If $f: (B_1, T_1) \rightarrow (B_2, T_2)$ be a mapping defined by $f(x') = (f(x))'$, $\forall x \in B_1$. Then f is fuzzy continuous.

Proof: Let $x \in B_1$ be any element. Let $v \in T_2$ be any open set. Then we show that $f^{-1}(v) \in T_1$. Now,

$$f^{-1}(v)(x) = v(f(x)) = v((f(x))') = v(f(x')) = f^{-1}(v)(x') \quad [\text{As } v \in T]$$

So, $f^{-1}(v) \in T$. Hence f is fuzzy continuous.

Theorem(4.4): Let the mapping $f: (B_1, T_1) \rightarrow (B_2, T_2)$ be fuzzy continuous. Then

$$f(x') = (f(x))' \text{ or } (f(x)), \quad \forall x \in B_1.$$

Proof: As f is continuous. Therefore, if v be any T_2 -open set, then $f^{-1}(v)$ is T_1 -open set.

Let $x \in B_1$ be any element, then we have $f^{-1}(v)(x) = f^{-1}(v)(x') \Rightarrow v(f(x)) = v(f(x'))$.

Let $H = \{f(x), (f(x))'\}$. Clearly, $H \subseteq B_2$. Define the fuzzy set θ on B_2 as follow:

$$\theta(y) = \begin{cases} \lambda & \text{if } y \in H \\ 0 & \text{if } y \notin H \end{cases} \quad \text{where } \lambda \in (0, 1]$$

Obviously, $\theta \in T_2$. As proved earlier, $\theta(f(x)) = \theta(f(x'))$. This means that either $f(x') = (f(x))'$ or $(f(x))$.

Hence proved.

Note(4.5): If f is one-one or $f(x') \neq (f(x))'$ for all $x \in B_1$, then we have the following theorem:

Theorem(4.6): Let $f: (B_1, T_1) \rightarrow (B_2, T_2)$ be a bijective mapping. Then f is fuzzy continuous if and only if $f(x') = (f(x))'$, $\forall x \in B_1$.

Theorem(4.7): Boolean algebraic fuzzy topological space is a fuzzy topological property.

Proof: Let $(B, +, \bullet, ', 0_B, 1_B)$ be a Boolean algebra and (B, T) be a BAFTS.

Suppose that $f: (B, T) \rightarrow (X, T')$ be a fuzzy homeomorphism. We will show that (X, T') is also a BAFTS. To prove this, we first show that X is a Boolean algebra under the operation:

For any $y_i \in X$ $\exists$'s unique $x_i \in B$ such that $f(x_i) = y_i$, where $i = 1, 2, 3$. We have

$$y_1 \otimes y_2 = f(x_1) \otimes f(x_2) = f(x_1 \bullet x_2) \text{ i.e., } y_1 \otimes y_2 = f(f^{-1}(y_1) \bullet f^{-1}(y_2)) = f(x_1 \bullet x_2) \text{ and}$$

$$y_1 \oplus y_2 = f(x_1) \oplus f(x_2) = f(x_1 + x_2) \text{ i.e., } y_1 \oplus y_2 = f(f^{-1}(y_1) + f^{-1}(y_2)) = f(x_1 + x_2).$$

Clearly, $y_1 \otimes y_2 = y_2 \otimes y_1$ and $y_1 \oplus y_2 = y_2 \oplus y_1$. Moreover, it is easy to show

$$\text{that } y_1 \otimes (y_2 \otimes y_3) = (y_1 \otimes y_2) \otimes y_3 \text{ and } y_1 \oplus (y_2 \oplus y_3) = (y_1 \oplus y_2) \oplus y_3.$$

$$\begin{aligned} \text{Also, } y_1 \otimes (y_2 \oplus y_3) &= y_1 \otimes [f(x_2 + x_3)] = f(x_1) \otimes [f(x_2 + x_3)] = f\{x_1 \bullet (x_2 + x_3)\} = f\{x_1 \bullet x_2 + x_1 \bullet x_3\} \\ &= f(x_1 \bullet x_2) \oplus f(x_1 \bullet x_3) = y_1 \otimes y_2 \oplus y_1 \otimes y_3. \end{aligned}$$

$$\text{Similarly, we can show that } y_1 \oplus (y_2 \otimes y_3) = (y_1 \oplus y_2) \otimes (y_1 \oplus y_3)$$

There exists elements $f(1_B) = 1_X$ and $f(0_B) = 0_X$ in X such that $y \otimes 1_X = y$ and $y \oplus 0_X = y$, $\forall y \in X$

Existence of inverse: let $y \in X$. Then $\exists x \in B$ such that $y = f(x)$.

We show that $y' = f(x')$ is the complement of y .

$$\text{Now, } y \otimes y' = f(x) \otimes f(x') = f(x \bullet x') = f(1_B) = 1_X \text{ and } y \oplus y' = f(x) \oplus f(x') = f(x + x') = f(0_B) = 0_X.$$

Hence X is a Boolean algebra under the operations $\oplus$, $\otimes$ and $'$.

Next, we show that the topology T' is the BAFT on X .

Let T'' be the BAFT on X . Then we show that $T' = T''$.

Let $y \in X$ be any element then $\exists$ a unique $x \in B$ such that $f(x) = y$.

$$\text{Let } v \in T'. \therefore f^{-1}(v) \in T.$$

Also, because T is a BAFT on B

$$\therefore f^{-1}(v)(x) = f^{-1}(v)(x'); \forall x \in B$$

$$\text{i.e., } v(f(x)) = v(f(x')) = v((f(x))')$$

$$v(y) = v(y'); \forall y \in X$$

$$v \in T, \quad T' \subseteq T'.$$

Conversely, let $v \in T''$. Then for any element $y \in X$, $\exists$ unique $x \in B$ such that $f(x) = y$.

$$\text{Now } v(y) = v(y') \Rightarrow v(f(x)) = v((f(x))') = v(f(x'))$$

$$\text{i.e., } f^{-1}(v)(x) = f^{-1}(v)(x') \Rightarrow f^{-1}(v) \in T.$$

As f is fuzzy open mapping $\therefore f(f^{-1}(v)) \in T'$ i.e. $v \in T'$.

Thus $v \in T' \Rightarrow v \in T$ so $T' \subseteq T$. Hence $T' = T$.

Corollary(4.8): Let $f: (B_1, T_1) \rightarrow (B_2, T_2)$ be fuzzy continuous mapping and $K = \{x \in B_1 : f(x) = 0_{B_2} \text{ and } f(x') = 1_{B_2}\}$. Then the topology induced on the set K is a BAFT.

Proof: Let $T_K = \{\mu \cap \chi_K : \forall \mu \in T_1\}$. Then we claim that T_K is a BAFT on K .

Let $x \in K$ be any element, then $f(x) = 0_{B_2}$ and $f(x') = 1_{B_2}$.

Let $v \in T_2$ be any open set in B_2 . As f is continuous $\therefore f^{-1}(v) \in T_1$.

$$\begin{aligned} \text{Now, } (f^{-1}(v) \cap \chi_K)(x) &= \min\{f^{-1}(v)(x), \chi_K(x)\} \\ &= \min\{v(f(x)), \chi_K(x)\} \\ &= \min\{v(0_{B_2}), \chi_K(x)\} \quad [\text{As } \chi_K(x) = 1] \\ &= v(0_{B_2}) = v(1_{B_2}) = v(f(x')) \\ &= \min\{v(f(x')), \chi_K(x')\} \quad [\text{As } \chi_K(x') = 1] \\ &= \min\{f^{-1}(v)(x'), \chi_K(x')\} \\ &= (f^{-1}(v) \cap \chi_K)(x') \end{aligned}$$

Hence T_K is a BAFT on K .

5. Product of Boolean Algebraic Fuzzy Topological Spaces

In this section, we study the product of Boolean algebraic fuzzy topological spaces. We also discuss similarities with difference from product of fuzzy topological spaces

Theorem (5.1): Let (B_1, T_1) and (B_2, T_2) be two BAFTS. Then product fuzzy topology on $B_1 \times B_2$ is contained in the BAFT on $B_1 \times B_2$. The equality between the two topology does not need to be hold.

Proof: We know that $B_1 \times B_2$ is a Boolean algebra under the operation defined by

$$\begin{aligned} (x_1, y_1) \otimes (x_2, y_2) &= (x_1 \bullet x_2, y_1 \bullet y_2); (x_1, y_1) \oplus (x_2, y_2) = (x_1 + x_2, y_1 + y_2); \forall (x_1, y_1), (x_2, y_2) \in B_1 \times B_2 \text{ and} \\ (x, y)' &= (x', y'), \forall (x, y) \in B_1 \times B_2. \end{aligned}$$

The element $(0_1, 0_2)$ and $(1_1, 1_2)$ be the lower bound and upper bound element of $B_1 \times B_2$. Let (x, y) be any element of $B_1 \times B_2$. Then (x', y') is the complement of (x, y) in $B_1 \times B_2$. Let T' be the product fuzzy topology on $B_1 \times B_2$ and T be the BAFT on $B_1 \times B_2$.

We show that $T' \subseteq T$. Let $\theta \in T'$ and let $z = (x, y)$ be any element of $B_1 \times B_2$.

Then $\theta(z) = \theta(x, y) = (\mu \times v)(x, y) = \min\{\mu(x), v(y)\} = \min\{\mu(x'), v(y')\} = (\mu \times v)(x, y)' = \theta(x, y)' = \theta(z')$.

so that $\theta \in T$. Hence $T' \subseteq T$.

Next, we show that the equality between T and T' does not need to behold.

Let $B_1 = (D_6, +, \cdot, ')$ and $B_2 = (D_{30}, +, \cdot, ')$. Let T_1, T_2 be the BAFT on B_1 and B_2 respectively.

Further let T' be the product fuzzy topology on $B_1 \times B_2$ and T be the BAFT on $B_1 \times B_2$.

Consider the subset $H = \{(2, 6), (3, 5)\}$ of $B_1 \times B_2$. Define the fuzzy set η on $B_1 \times B_2$ such that $\eta((x, y)) = \lambda$

If $(x, y) \in H$ and 0 otherwise, where $\lambda \in (0, 1]$.

Clearly, $\eta(x, y)' = \eta(x, y); \forall (x, y) \in B_1 \times B_2$ and so, $\eta \in T$.

But the smallest T' -open set containing η is $\mu \times v$, where $\mu \in T_1$ and $v \in T_2$ defined as:

$$\mu(x) = \begin{cases} \lambda, & \text{if } x = 2, 3 \\ 0, & \text{otherwise} \end{cases} \quad v(x) = \begin{cases} \lambda, & \text{if } x = 6, 7 \\ 0, & \text{otherwise} \end{cases}$$

$$(\mu \times v)(x, y) = \min\{\mu(x), v(y)\} = \lambda; \text{ if } (x, y) \in \{(2, 6), (2, 5), (3, 6), (3, 5)\} = \{2, 3\} \times \{6, 5\}$$

$$\text{And } (\mu \times v)(x, y) = 0; \text{ if } (x, y) \notin \{2, 3\} \times \{6, 5\}.$$

Thus, $\eta \notin T'$. Hence, $T' \neq T$.

Remark(5.2): In the above theorem(5.1), if we have $B_1 = B_2 = B$ (say), even then the product fuzzy topology on $B \times B$ is not same as the BAFT on $B \times B$.

6. Conclusion

In this paper, the notion of Boolean algebraic fuzzy topological spaces is introduced. It has been observed that the Boolean algebraic fuzzy topology is different from discrete and in discrete fuzzy topology, the cases when they have been examined. It is seen that Boolean algebraic fuzzy topology is a hereditary property subject to the Boolean subalgebra of the Boolean algebra B . Also it is noticed that Boolean algebraic fuzzy topology is a topological and productive property.

References

- [1] C.L.Chang, “*Fuzzy Topological Spaces*”, J.Math.Anal.Appl., 24(1968), 182–190.
- [2] R. Lowen, “*Fuzzy Topological Spaces and Fuzzy Compactness*”, J. Math. Anal. Appl., 56 (1976), 621–633.
- [3] A. Parvathi, K.N. Meenakshi, “*Fuzzy Topological Boolean Algebras*”, Indian J. pure appl. Math. 28(12), (1997), 1639-1648.
- [4] A.Rosenfeld, “*Fuzzy Groups*”, J.Math.Anal.Appl., 35(1971), 512–517.
- [5] Shaoquan Sun, “*Generalized Fuzzy Subalgebras of Boolean Algebras*”, Proceedings of

the 2009 International Workshop on Information Security and Application (IWISA 2009) Qingdao, China, November 21-22, (2009).

- [6] C.K.Wong, “*Fuzzy Topology, Products and Quotient Theorems*”, J.Math.Anal.Appl., 45(1974), 512–521.
- [7] Shaoquan Sun, “*Fuzzy subalgebras and Fuzzy ideals of Boolean algebra*”, Fuzzy Systems and Mathematics, vol. 20(1) (2006) pp. 90-94.
- [8] L.A.Zadeh, “*Fuzzy sets, Information and Control*”, 8(1965), 338–353.
- [9] Elliott Mendelson, “*Schaum's outline of Boolean Algebra and Switching Circuits*”, McGrawHill, (1970), ISBN 07-041460-2.
- [10] N.Palaniappan, “*Fuzzy Topology*”, 2nd Edition Narosa Publishing House, India.

